

# Mathematical Modelling Exam

June 7th, 2023

This is an open book exam. You are allowed to use your notes, books and any other literature. You are NOT allowed to use any communication device. You have 90 minutes to solve the problems.

1. **[15 points]** Let

$$A = \begin{bmatrix} 5 & -1 \\ 1 & -5 \\ 5 & -1 \\ 1 & -5 \end{bmatrix} \quad \text{and} \quad c = \begin{bmatrix} 1 \\ -1 \\ 0 \\ 0 \end{bmatrix} \quad (1)$$

be a matrix and a vector. The singular value decomposition (SVD) of  $A$  is equal to

$$U\Sigma V^T = \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 6\sqrt{2} & 0 \\ 0 & 4\sqrt{2} \end{bmatrix} \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}^T.$$

- (a) Compute the lengths  $\alpha_1, \alpha_2$  of both projections of the vector  $c$  to the left singular vectors of  $A$  and compute the projection  $p$  of  $c$  to the orthogonal complement of the span of the left singular vectors of  $A$ .
- (b) In the notation above show that for any matrix  $A \in \mathbb{R}^{4 \times 2}$  it holds that

$$\begin{bmatrix} A & c \end{bmatrix} = \begin{bmatrix} U & \frac{p}{\|p\|} \end{bmatrix} \begin{bmatrix} \Sigma & \begin{array}{c} \alpha_1 \\ \alpha_2 \end{array} \\ 0_{1 \times 2} & \|p\| \end{bmatrix} \begin{bmatrix} V & 0_{2 \times 1} \\ 0_{1 \times 2} & 1 \end{bmatrix}^T. \quad (2)$$

Here,  $\|p\|$  stands for the usual Euclidean norm of the vector  $p$  and  $0_{i \times j}$  stands for the  $i \times j$  matrix with zero entries.

- (c) Using (1b) for the given matrix  $A$  and the vector  $c$  from (1), compute the singular values of the matrix  $\begin{bmatrix} A & c \end{bmatrix}$ .

Hint: Observe that  $\begin{bmatrix} U & \frac{p}{\|p\|} \end{bmatrix}$  and  $\begin{bmatrix} V & 0_{2 \times 1} \\ 0_{1 \times 2} & 1 \end{bmatrix}$  are orthogonal matrices and hence you only need to compute the singular values of the middle matrix in the factorization (2). For all points give a short explanation why this last conclusion holds.

2. **[10 points]** Let  $D = \text{diag}(d_1, \dots, d_n)$  be a diagonal matrix with  $0 < d_1 < \dots < d_n$  and let  $z = \begin{bmatrix} z_1 & z_2 & \dots & z_n \end{bmatrix}^T \in \mathbb{R}^n$  be a vector. We form a matrix  $M = \begin{bmatrix} D & z \end{bmatrix}$ , i.e.,

$$M = \begin{bmatrix} d_1 & & & z_1 \\ & d_2 & & z_2 \\ & & \ddots & \vdots \\ & & & d_n & z_n \end{bmatrix}$$

It turns out that nonzero singular values  $\sigma_1, \dots, \sigma_n$  of the matrix  $M$  are solutions of the nonlinear equation

$$1 + \sum_{i=1}^n \frac{z_i^2}{d_i^2 - w^2} = 0$$

and they satisfy the interlacing property

$$0 < d_1 < w_1 < d_2 < w_2 < \dots < d_n < w_n < d_n + \|z\|.$$

Let  $n = 3$ ,  $d_1 = 1$ ,  $d_2 = 2$ ,  $d_3 = 3$  and  $z_1 = z_2 = z_3 = 1$ .

**Task:** Perform one step of Newton's method with a suitably chosen initial approximation  $w^{(0)}$  to estimate the smallest and the largest singular values  $\sigma_1, \sigma_3$ , i.e., repeat the procedure twice with different, meaningfully chosen initial approximations.

3. **[15 points]** Let  $f(u, v) = (f_1(u, v), f_2(u, v), f_3(u, v))$ ,  $u_1 \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ ,  $u_2 \in [0, 2\pi)$  be a parametrization of the torus:

$$f(u_1, u_2) = ((2 + \cos(u_1)) \cos(u_2), (2 + \cos(u_1)) \sin(u_2), \sin(u_1)).$$

- (a) Compute the matrix  $G$ , called the metric tensor, and its inverse  $G^{-1}$ :

$$G = \begin{bmatrix} \sum_{k=1}^3 \frac{\partial f_k}{\partial u_1} \frac{\partial f_k}{\partial u_1} & \sum_{k=1}^3 \frac{\partial f_k}{\partial u_1} \frac{\partial f_k}{\partial u_2} \\ \sum_{k=1}^3 \frac{\partial f_k}{\partial u_2} \frac{\partial f_k}{\partial u_1} & \sum_{k=1}^3 \frac{\partial f_k}{\partial u_2} \frac{\partial f_k}{\partial u_2} \end{bmatrix}, \quad G^{-1} = \begin{bmatrix} h_{11} & h_{12} \\ h_{21} & h_{22} \end{bmatrix}.$$

- (b) For  $i, j, k = 1, 2$ , the so-called Christoffel symbols  $\Gamma_{ij}^k$  are defined by:

$$\Gamma_{ij}^k = \sum_{\ell=1}^2 \left\langle \begin{bmatrix} \frac{\partial^2 f_1}{\partial u_i \partial u_j} \\ \frac{\partial^2 f_2}{\partial u_i \partial u_j} \\ \frac{\partial^2 f_3}{\partial u_i \partial u_j} \end{bmatrix}, \begin{bmatrix} \frac{\partial f_1}{\partial u_\ell} \\ \frac{\partial f_2}{\partial u_\ell} \\ \frac{\partial f_3}{\partial u_\ell} \end{bmatrix} \right\rangle h_{\ell k},$$

where  $\langle \cdot, \cdot \rangle$  stands for the usual inner product of vectors. Compute  $\Gamma_{11}^1$ .

- (c) The shortest paths on torus are obtained by solving the following second order system of differential equations:

$$\begin{aligned} \frac{d^2 u_1}{dt^2} + \sum_{i,j=1}^2 \Gamma_{ij}^1 \frac{du_i}{dt} \frac{du_j}{dt} &= 0, \\ \frac{d^2 u_2}{dt^2} + \sum_{i,j=1}^2 \Gamma_{ij}^2 \frac{du_i}{dt} \frac{du_j}{dt} &= 0. \end{aligned} \tag{3}$$

Except  $\Gamma_{11}^1$  you computed in (3b), the remaining Christoffel symbols are

$$\begin{aligned} \Gamma_{11}^2 &= 0, \quad \Gamma_{12}^1 = \Gamma_{21}^1 = 0, \quad \Gamma_{12}^2 = \Gamma_{21}^2 = -\frac{\sin(u_1)}{2 + \cos(u_1)}, \\ \Gamma_{22}^1 &= (2 + \cos(u_1)) \sin(u_2), \quad \Gamma_{22}^2 = 0. \end{aligned}$$

Write down the system (3) explicitly.

- (d) Translate the system from (3c) to the first order system of differential equations by introducing new variables

$$x_1(t) = u_1(t), \quad x_2(t) = \frac{du_1}{dt}, \quad x_3(t) = u_2(t), \quad x_4(t) = \frac{du_2}{dt}.$$

- (e) For the initial point  $(u_1(0), u_2(0)) = (0, 0)$  and derivatives  $(\frac{du_1}{dt}(0), \frac{du_2}{dt}(0)) = (1, 1)$  perform one step of Euler's method on the system from (3d) with a step size  $h = 0.1$ .